

Safflower XVI-3

Chemical Control for Safflower

Gary L. Hein

Insecticide	Product per Acre	Preharvest Interval, Remarks
For control of Lepidoptera larvae		
Bacillus thuringiensis var. kurstaki (several products)	See labels for rates.	PHI 0 days; REI 4 hrs.
For control of Lygus bug and thrips		
Supracide 25WP ^R , 2E ^R	25WP: 2.0 lbs/A 2E: 2.0 lbs./A	PHI 28 days; REI 48 hrs.
For control of aphids and mites		
Trilogy	See label for rates.	Do not apply when bees are active. REI 4 hrs.
Seed Treatments for flea beetle, wireworm control		
Cruiser 5FS	10.24 oz/100 lb seed	Treated by seed supplier only.

^RRestricted use pesticide ¹Labelled for chemigation.

The information herein is supplied with the understanding that no discrimination is intended and that listing of commercial products, necessary to this guide, implies no endorsement by the authors or the Extension Services of Nebraska, Colorado, Wyoming or Montana. Criticism of products or equipment not listed is neither implied nor intended. Due to constantly changing labels, laws and regulations, the Extension Services can assume no liability for the suggested use of chemicals contained herein. Pesticides must be applied legally complying with all label directions and precautions on the pesticide container and any supplemental labeling and rules of state and federal pesticide regulatory agencies. State rules and regulations and special pesticide use allowances may vary from state to state: contact your State Department of Agriculture for the rules, regulations and allowances applicable in your state and locality.

Categories: Safflower, Chemical, Armyworm, Loopers, Grasshoppers, Lygus Bugs, Thrips, Insects

Date: 04/19/2006